

The state versus amyloid- β : the trial of the most wanted criminal in Alzheimer disease

Catherine A. Rottkamp^a, Craig S. Atwood^a, James A. Joseph^b, Akihiko Nunomura^c,
George Perry^a, Mark A. Smith^{a,*}

^a *Institute of Pathology, Case Western Reserve University, 2085 Adelbert Road, Cleveland, OH 44106, USA*

^b *Human Nutrition Research Center on Aging, Tufts University, Boston, MA, USA*

^c *Department of Psychiatry and Neurology, Asahikawa Medical College, Asahikawa, Japan*

Received 2 May 2001; accepted 13 June 2001

Abstract

Investigators studying the primary culprit responsible for Alzheimer disease have, for the past two decades, primarily focused on amyloid- β (A β). Here, we put A β on trial and review evidence amassed by the prosecution that implicate A β and also consider arguments and evidence gathered by the defense team who are convinced of the innocence of their client. As in all trials, the arguments provided by the prosecution and defense revolve around the same evidence, with opposing interpretations. Below, we present a brief synopsis of the trial for you, the jury, to decide the verdict. **Amyloid- β : guilty or not-guilty?**

© 2002 Elsevier Science Inc. All rights reserved.

Keywords: Alzheimer disease; Cell death; Neurodegeneration; Neurotoxicity; Oxidative stress; Phosphorylated tau

The execution date for amyloid- β (A β) is now imminent. Vaccination of A β to Alzheimer disease (AD) patients, via an intranasal or intravenous route, is, in many people's opinion, going to improve the disease course by removing (killing) intracerebral A β senile plaques [112]. However, many others argue that a gross miscarriage of justice is about to occur and that executing A β still leaves the killer loose in the brain. The aim of this review is to, in effect, put A β on trial in front of a jury of readers by presenting evidence used by the prosecutors and defenders of A β . The prosecution, backed by public opinion eager to pin the blame on something, argues for guilt based on the fact that amyloid burden in the brain is correlated with dementia, that A β deposits are found in regions of the brain susceptible to neurodegenerative processes and that A β production is increased in all inherited forms of the disease [114]. Additionally, A β is a known killer of neurons in vitro [95,145] and surely must wreak the same carnage in the human brain. On the other hand, the defense, in the face of an overwhelming number of publications implicating A β as the major culprit in AD pathophysiology, argue for the innocence of A β as either an innocent bystander or a maligned protector of the brain [125]. The following represents a brief synopsis of the trial

and, while the scientific debates continue to encircle this issue, it is our hope that this trial will define the questions that remain to be answered by both sides.

1. The Prosecution

When investigating the crime scene of AD, namely the brain tissue of patients, it is quite obvious why A β is the primary suspect in disease pathogenesis. In fact, some of the earliest detectives studying the biochemistry of the AD brain determined that the A β peptide is the major constituent in two of the most distinctive pathologies, namely senile plaques and cerebral amyloid angiopathy [39,40,75]. A β is derived by proteolytic cleavage of A β precursor protein (A β PP), a protein of unestablished cellular function that has the general motif of a surface receptor [61,87]. Regardless of the fact that A β PP is a transmembrane protein in the neuronal plasma membrane, a secretory pathway in the Golgi apparatus processes the large majority of the protein before it ever reaches the cell surface [19,24,67]. While it was originally thought that A β represents an abnormal cleavage product, A β has been since been established as a normal product of neuronal A β PP metabolism, found in the cerebral spinal fluid (CSF) and serum of healthy individuals [45,118]. Differential activity between three different

* Corresponding author. Tel.: +1-216-368-3670; fax: +1-216-368-8964.
E-mail address: mas21@po.cwru.edu (M.A. Smith).

secretases, α , β and γ , at their specific cleavage sites yields a number of different products, including $A\beta_{1-40}$ and $A\beta_{1-42}$ [120]. While $A\beta_{1-40}$ is the predominant product of this proteolytic pathway, $A\beta_{1-42}$ is far more fibrillogenic in vitro and is the major $A\beta$ species present in the core of senile plaques (both AD and non-AD related) [13,56]. The deposition of $A\beta_{1-40}$ and $A\beta_{1-42}$ into senile plaques likely begins with the nucleation of soluble $A\beta_{1-42}$ into fibrils followed by accumulation of normally soluble $A\beta_{1-40}$ [56]. Micro-environmental changes in the brain, such as pH, metal ion availability and oxidants, likely impact upon $A\beta$ structural conformation and its deposition in amyloid plaques [6,123]. As of late a great deal of attention has also been focused on the fact that soluble forms of amyloid that are pre-fibrillar may also be involved in AD pathogenesis [69]. However, soluble forms of $A\beta$, including oligomers, correlate quantitatively with the number of senile plaques [130] and, therefore, all of the prosecution's (and defense's) arguments apply equally to all forms of $A\beta$.

The destructive nature of $A\beta$ is evident from a close examination of histological preparations in the immediate vicinity of senile plaques revealing degenerative dendritic processes surrounding and infiltrating the plaques [36]. Additionally, regions severely affected by disease, including the hippocampus and frontotemporal cortices, show colocalization between $A\beta$ plaques and neuronal cell death [107]. This led investigators to explore whether $A\beta$ is toxic to neurons in both in vitro culture assays and in the intact brain of animals. The results of these experiments at first seemed extremely contradictory, stemming from lot-to-lot variability in the peptide and the lack of proper control over whether $A\beta$ was aggregated into fibrils of β -sheet conformation [28]. Nonetheless, it is now established that fibrillation of $A\beta$ is required to obtain neurotoxic effects [71,97,146] and that it is inherently toxic to neurons and clonal cell lines in culture [95,145]. Toxicity of the peptide has been shown to reside in between amino acids 25 and 35 [98]. The neurotoxicity of the peptide in vivo was likewise assessed by infusion of the peptide in a variety of animal models. Notably, intracortical injection of $A\beta_{1-42}$ or $A\beta_{25-35}$ fragments into aged rats or primates produced lesions reminiscent to those seen in AD patients [66]. In vivo injection of $A\beta$ into the brain also led to neurodegeneration and dystrophic neurites in hippocampal neurons [95,96].

The source of $A\beta$ toxicity has yet to be established, however, a number of theories have been advanced. The possibility that $A\beta$ may act through a cell surface receptor and thereby trigger an intracellular signaling cascade, while not being eliminated as a possibility, is supported by little experimental evidence. Numerous studies have since supported the idea that an oxidative event is critical for $A\beta$ toxicity (reviewed in [76]). It is thought that the peptide is capable of generating reactive oxygen species (ROS), which is supported by the fact that $A\beta$ peptides can induce the generation of H_2O_2 [48,54,55], can stimulate inflammatory cells to produce ROS [1,15,81,135] and that neurotoxicity

can be attenuated by administration of anti-oxidants and free radical scavengers, such as Vitamin E [9]. The prooxidant potential of $A\beta$ also is supported by in vivo evidence where deposits are associated with oxidative damage [7,122,124] and such damage is, like $A\beta$ deposition [113], viewed as an extremely proximal event in disease pathogenesis [89,90]. However, while it is clear that $A\beta$, either directly or indirectly, promotes oxidative stress and that toxicity can be attenuated by anti-oxidants, the precise mechanism by which amyloid deposition leads to increased oxidative stress remains elusive. Indeed, while studies have suggested that the neurotoxicity of aggregated $A\beta$ is mediated by its ability to induce oxidative stress via the spontaneous generation of free radicals and ROS [48], this proposition has been questioned on theoretical and methodological grounds [31,110,133]. Instead, it now appears that the oxidant effects of $A\beta$ are mediated by its interaction with redox-active metals such as iron and copper since chelation treatment of $A\beta$ significantly attenuates toxicity [109]. Significantly, $A\beta$ has an unusually high affinity for both iron and copper [6,29,53] and is capable of reducing these metals with subsequent production of hydrogen peroxide and oxidized amyloid [54,55]. The relevance of this mechanism to disease pathogenesis is highlighted by the association of redox active metals with senile plaques in AD [111,123]. In addition, the deposition of this normally soluble cellular protein promotes a chronic inflammatory response of the AD brain, whereby activated microglia release ROS as part of the respiratory burst (reviewed in [8]). Therefore, $A\beta$ likely has much to answer for with regards to the oxidative damage observed in the AD brain.

Perhaps the strongest evidence for the crucial role played by $A\beta$ in AD pathogenesis has been the characterization of the mutations that underlie familial early onset cases of the disease. All of these inherited mutations directly or indirectly affect the processing and accumulation of $A\beta$. The most straightforward form of familial Alzheimer disease (FAD) is caused by point mutations in $A\beta$ PP in regions that are involved in the proteolytic processing of the peptide [41,70]. It is thought that these mutations accelerate the onset of AD into the fourth decade by increasing the ratio of $A\beta_{1-42}/A\beta_{1-40}$, thereby increasing the relative amount of the more fibrillogenic form [128]. A double mutation at positions 670/671 (Swedish mutation) increases the production of total $A\beta$ and thereby increases the load of $A\beta_{1-42}$ without changing the relative ratio [18,23]. The fact that an increase in total $A\beta$ load can accelerate the deposition of $A\beta$ is supported by the neuropathology demonstrated in patients with Down syndrome, a disorder caused by trisomy of chromosome 21, where the $A\beta$ PP gene is localized. It is thought that the overexpression of $A\beta$ PP in these individuals [44] causes the formation of $A\beta$ plaques very similar to those seen in the AD brain. The most common form of FAD is caused by mutations in one of the two presenilin genes (PS1 on chromosome 14 or PS2 on chromosome 1) (reviewed in [43]). While the function of the presenilin genes

are not completely elucidated, it is known that the presenilin proteins localize to the endoplasmic reticulum and Golgi apparatus [65] and form stable complexes with A β PP [142]. Most importantly however, missense mutations in the presenilin genes likewise increase the ratio of A β _{1–42}/A β _{1–40} [25,131]. Finally, it has been demonstrated that one allele of the apolipoprotein E gene, namely ApoE4 predisposes individuals to the development of late-onset AD [26]. Of the three alleles (also including ApoE2 and ApoE3), ApoE4 has the greatest affinity for A β , is found associated with senile plaques and is thought to accelerate fibrillogenesis [141]. Interestingly, ApoE2 is capable of inhibiting the formation of fibrils and is protective against the development of AD [27].

This strong evidence implicating A β in AD pathogenesis led to the supposition that generation of transgenic animals either over expressing A β PP or containing a mutation in A β PP that affects processing of the full length protein thereby leading to an increase in the A β _{1–42}/A β _{1–40} ratio, may mimic the symptomatology of AD. Taken as a group, the various transgenic mice strains that have been produced have demonstrated that overexpression of A β PP or overproduction of the A β _{1–42} peptide fragment is sufficient to cause deposition of the peptide into senile plaque-like structures and may in fact be responsible for at least some of the neurodegeneration in AD (reviewed in [50]). Indeed, despite the fact that each of the different constructs introduced yielded somewhat different phenotypes, some aspect of AD pathophysiology are apparent in each. For example, Games et al. created a transgenic mouse expressing human A β PP with the Val717Phe mutation at 10 times the endogenous level and these animals developed amyloid plaques in the hippocampus, cerebral cortex and corpus collosum by 6–9 months of age and also show synaptic loss and astrogliosis, although they show no behavioral abnormalities [35]. Among the first models to demonstrate behavioral changes reminiscent of those seen in AD, overexpressed A β PP containing the Swedish double mutation (Lys670Asn/Met671Leu). In addition to a marked increase in levels of A β in the CSF and deposition of amyloid plaques, these mice demonstrated marked deficits in spatial learning, as demonstrated by Morris water maze, by the age of 9 months [52]. Although no neurotoxicity was observed in these mice, it is thought that their impaired spatial learning, which is correlated with long-term potentiation (a model for memory), is related to synaptic loss. Interestingly, these mice also displayed oxidative stress in association with the plaques, much like that seen in AD [124]. Neurotoxicity has also been seen accompanied by an increased mortality rate, with 50% of mice dying by 12 months of age compared to an average of 24 months in controls, in mice overexpressing A β PP [68]. Although there are differences in the details of these studies, it is clear that A β can independently cause AD-related pathology and some behavioral defects.

Taken together, the aforementioned evidence clearly points the suspicion of doubt at A β as being an instrumental, if not the sole, culprit for causing disease.

2. The Defense

It is obvious that proponents of the causative role of A β in AD have a great deal of evidence in their arsenal and the preceding discussion would be sufficient to convince most juries of its guilt. Unfortunately, however, much of this evidence is circumstantial and in many cases the prosecution failed to discuss the caveats that create uncertainties as to the role of A β in AD. While we by no means intend to suggest that A β plays a negligible role in disease pathogenesis, it is very unlikely that the peptide is the main mediator of neurodegeneration.

The first, and perhaps most poignant, argument is the fact that deposition of A β into senile plaques is by no means specific to AD patients and in fact seems to be characteristic of normal aging [30]. The incidence of amyloid plaques in control individuals increases with age, as does the incidence of AD, and the number of plaques in cognitively normal individuals can rival those seen in advanced disease [74]. Even within those patients with AD, there is only a weak correlation between the burden of A β and neuronal loss or cognitive impairment [30,86]. Additionally, increased amyloid production and deposition is observed in response to injury to the central nervous system, including ischemia and head trauma [37,38,106]. Also, despite marked amyloid deposition in the brains of Down syndrome patients by the fourth decade, there is little evidence of further cognitive decline. Therefore, it appears that A β deposition is insufficient to develop full fledged AD. Rather it seems that A β may be produced in the disease as a cellular protective factor to compensate for the primary insult that causes AD [58,59,125]. This is not to imply that, in attempting to respond to cellular stresses, A β does not produce some cellular destruction while maintaining the integrity of the whole. However, it does require an underlying pre-existing stress, i.e. the presumable causative insult of AD.

A review of the literature would indicate that the underlying stress is an energetic one, since a shortage of energy supply (and Ca(II) overload) induces an upregulation of A β PP expression. Ischemia, hypoglycemia and traumatic brain injury, a condition that has been shown to put neurons under metabolic stress [143], all upregulate A β PP and its mRNA in animal models and culture systems [46,57,83,115–117,147]. Not only does energy shortage and Ca(II) dysregulation promote A β PP expression, but they also route the metabolism of A β PP from the non-amyloidogenic to the amyloidogenic pathway. Inhibition of mitochondrial energy metabolism alters the processing of A β PP to generate amyloidogenic derivatives [34,77], while oxidative stress has been shown to increase the generation of A β [32,82,93]. Consistent with this response A β has been detected in the human brain several days after traumatic brain injury [38]. This fits well with the role of A β PP as an acute phase reactant upregulated in neurons, astrocytes and microglial cells in response to inflammation and a multitude of associated cellular stresses including

axonal injury [11,38], loss of innervation [138], excitotoxic stress [92,132], heat shock [22], oxidative stress [32,144], aging [49,88,134] and inflammatory processes [12]. Other pro-inflammatory stimuli that mediate the synthesis and release of A β PP include IL-1 β [16,42] and TNF α converting enzyme [17]. The increased expression of A β PP by these stress conditions is likely a result of decreased energy supply.

The increased generation of A β under conditions of energetic stress may therefore be a response to the oxidative challenge observed in the brain in AD and following injury. In this vein, A β may in fact play a protective role. In support of this claim, the A β amyloid burden of the AD-affected brain has been shown to be significantly negatively correlated with oxidative stress markers [29,89,90] and in situ, soluble A β levels are inversely correlated with synaptic loss [72]. This argues against the neurotoxic role of A β in vivo, as does the fact that cultured neurons can be cultured directly on top of isolated A β plaques with out any notable toxicity [20]. Some have suggested that the in vitro toxicity that has been sporadically shown in culture, and very unreliable in animal models, may in fact be an artifact of culture and not intrinsic to the peptide itself [109]. Notably, A β appears to blunt oxidative stress in vivo [90,91] likely acting as an anti-oxidant [14]. Moreover, nanomolar concentrations of A β can block neuronal apoptosis following trophic factor withdrawal [21]. These findings are consistent with the trophic and neuroprotective action of A β at physiological concentrations in deprived conditions and neonatal cells that have been reported during the last decade [9,60,64,73,100,119,127,129,139,140,145]. A β also has been shown to protect neurons from death following injection of saline or iron [10] and protect lipoproteins from oxidation in cerebrospinal fluid and plasma (the mechanism of which is thought to involve metal ion sequestration) [6,62,63]. Moreover, Andorn and Kalaria [2] recently showed that low concentrations of A β possess significant anti-oxidant activity in an ascorbate-stimulated-lipid-peroxidation assay of post-mortem human brain membrane preparations. Together, these data provide a plausible physiological explanation for the increased generation of A β in AD and following head trauma, one that is aimed at reducing oxidative damage thereby preventing ROS-mediated neuronal apoptosis [105] and promoting neurite outgrowth.

Neurons are particularly vulnerable to oxidative stress as a consequence of high oxygen utilization, the high unsaturated lipid content of neuronal membranes and the post-mitotic nature of primary neurons. As a result, the balance between the production of ROS and anti-oxidant defenses are essential for proper neuronal function. Normally, anti-oxidant defense systems prevent any damage potentially produced by oxygen radicals. However, in cases of age related neurodegeneration, there is considerable oxidative imbalance, as demonstrated by accumulation of reversible and permanent alteration of cellular proteins and

nucleic acids (reviewed in [108]). Given the fact that A β has been associated with the production of free radicals, it is essential to consider the spatio-temporal relationship between oxidative stress phenomena and A β deposition. Notably, oxidative stress is found in normal appearing susceptible neurons in AD brain and seems to be inversely correlated with A β deposition [89,90]. It therefore seems unlikely that A β accumulation is sufficient to explain the oxidative imbalance. Vulnerable neurons also display cell cycle abnormalities that are uncharacteristic of terminally differentiated neuronal populations. Despite the fact that successful nuclear division has yet to be demonstrated, a large number cell cycle regulators, both drivers and inhibitors of the cell cycle checkpoints, have been found in association with AD lesions and normal-appearing susceptible neurons [3–5,47,78–80,84,85,101–104,121,136,137,148–151]. Although the precipitating factors for this type of regulation have yet to be elucidated, a number of other AD-related phenomena can be linked to cell cycle abnormalities, including tau hyperphosphorylation, increased production and processing of A β PP and oxidative stress (reviewed in [104,108]).

Perhaps the strongest evidence for the role of A β in AD is that each of the familial forms of the disease involve a mutation or polymorphism of genes that are directly involved in A β PP processing (see above). A tremendous amount of effort and resources have been dedicated in the past several decades to determining the mechanism of disease of these mutations. While this line of research has been fruitful in characterizing FAD, it has proven only marginally useful to our understanding of sporadic AD, which by far represents the majority of cases. For example, mutations in A β PP have been identified in only 20–30 families world wide and represent less than 0.1% of the 15 million known cases of AD. Mutations in both presenilin 1 and 2, which are the most common genetic determinant of AD, only contribute an additional 120–130 families. While it is clear that mutations in these proteins involved in A β PP processing are capable of inducing amyloid neuropathies and dementia, no aberrant neurologies are observed usually for many decades, and even then, this is likely a result of the advanced deposition of A β in these individuals (the joint result of increased A β concentration and microenvironmental conditions) leading to the chronic neuroinflammation associated with the disease.

The prosecution's claim that the positive correlation between ApoE4 genotype and incidence of AD supporting a causative role for A β is likewise flawed. While it is true that ApoE4 has the greatest affinity for A β , is found associated with senile plaques and is thought to accelerate fibrillogenesis [141], this is not the sole or even the major physiological role of ApoE proteins. In the periphery, it is known that ApoE helps to regulate the transport and metabolism of lipids. It is also well established that the level of ApoE is elevated in response to injury in the peripheral and central nervous system [51] and, like A β , ApoE may thereby serve a protective role after ischemia or traumatic

brain injury by distributing phospholipids and cholesterol to injured neurons [99]. ApoE may also protect against oxidative injury and prevent the accumulation of lipid peroxidation end products, such as hydroxynonenal, which are prominent features in AD and acute brain injury. In this vein, several studies showed that patients with homozygous ApoE4 genotype have longer periods of unconsciousness and higher incidence of post-traumatic coma following severe traumatic brain injury [33,126]. In short, ApoE4 predisposes patients for any number of neurodegenerative processes and is another factor that is completely unspecific to AD. Therefore, much like in acute injury, ApoE4 may be associated with a higher incidence of AD because it is less efficient in protecting neurons from the causative insult and therefore may have very little to do with its affinity for A β .

In sum, the defense argues that A β simply represents an innocent bystander rather than the culprit of disease [94,125].

3. Note added in press

Since the article was originally written, the removal of A β from patients with AD has been attempted as a therapeutic strategy. The technique involved immunizing patients with A β , a strategy which showed promise in reducing plaque burden in transgenic mice [112]. However, the clinical trials were suspended in March of 2002 when many patients developed acute exacerbation of symptoms, including confusion and inability to perform tasks [153–155]. While Elan Pharmaceutical Corporation has yet to release the details of the trials [152], one wonders whether this is major victory for the defense who said all along that A β was a much maligned protector of the brain and that removal would only serve to increase injury [58,59,94]. However, unfortunately, there are no rules against double jeopardy in science and A β will surely be accused and brought to trial again.

4. Concluding remarks

AD is a devastating condition and patients, caregivers, clinicians and scientists are eager to find the culprit and cure the disease. Whether A β is the culprit, as argued by the prosecution, or a bystander, as argued by the defense, is clearly important to decipher. All investigations start with multiple suspects, even multiple prime suspects, and it is the hope of all involved that we have the answer soon and can put this killer to rest.

References

[1] Akama KT, Albanese C, Pestell RG, Van Eldik LJ. Amyloid-beta peptide stimulates nitric oxide production in astrocytes through an NF κ B-dependent mechanism. *Proc Natl Acad Sci USA* 1998;95:5795–800.

[2] Andorn AC, Kalaria RN. Factors affecting pro- and anti-oxidant properties of fragments of the β -protein precursor (β PP): implication for Alzheimer's disease. *J Alzheimer's Dis* 2000;2: 69–78.

[3] Arendt T, Holzer M, Grossmann A, Zedlick D, Bruckner MK. Increased expression and subcellular translocation of the nitrogen activated protein kinase kinase and mitogen-activated protein kinase in Alzheimer's disease. *Neuroscience* 1995;68:5–18.

[4] Arendt T, Rodel L, Gartner I, Holzer M. Expression of the cyclin-dependent kinase inhibitor p16 in Alzheimer's disease. *Neuroreport* 1996;7:3047–9.

[5] Arendt T, Holzer M, Gartner U. Neuronal expression of cyclin dependent kinase inhibitors of the INK4 family in Alzheimer's disease. *J Neural Transmission* 1998;105:949–60.

[6] Atwood CS, Huang X, Moir RD, Bacarra NM, Romano D, Tanzi RE, et al. Dramatic aggregation of Alzheimer A β by Cu(II) is induced by conditions representing physiological acidosis. *J Biol Chem* 1998;273:12817–26.

[7] Atwood CS, Smith MA, Perry G. Senile plaque composition and posttranslational modification of amyloid- β peptide and associated proteins. *Peptides* 2001;26:1339–46.

[8] Atwood CS, Huang X, Moir RD, Smith MA, Tanzi RE, Roher AE, et al. Neuroinflammatory responses in the Alzheimer's disease brain promote the oxidative post-translational modification of amyloid deposits. In: Iqbal K, Sisodia SS, Winblad B, editors. *Alzheimer's disease: advances in etiology, pathogenesis and therapeutics*. Chichester, UK: Wiley, 2001. p. 341–61.

[9] Behl C, Davis J, Cole GM, Shubert D. Vitamin E protects nerve cells from amyloid- β protein toxicity. *Biochem Biophys Res Commun* 1992;186:944–50.

[10] Bishop GM, Robinson SR. β -Amyloid helps to protect neurons from oxidative stress. *Neurobiol Aging* 2000;21:S226.

[11] Blumergs PC, Scott G, Manavis J, Wainwright H, Simpson DA, McLean AJ. Topography of axonal injury as defined by amyloid precursor protein and the sector scoring method in mild and severe closed head injury. *J Neurotrauma* 1995;12:565–72.

[12] Brugg B, Dubreuil YL, Huber G, Wollman EE, Delhay-Bouchaud N, Mariani J. Inflammatory processes induce beta-amyloid precursor protein changes in mouse brain. *Proc Natl Acad Sci USA* 1995;92:3032–5.

[13] Burdick D, Soreghan B, Kwon M, Kosmoski J, Knauer M, Henschen A, et al. Assembly and aggregation properties of synthetic Alzheimer's A4/beta amyloid peptide analogs. *J Biol Chem* 1992;267:546–54.

[14] Bush AI, Lynch T, Cherny RC, Atwood CS, Goldstein LE, Moir RD, et al. Alzheimer A β functions as a superoxide anti-oxidant in vitro and in vivo. *Soc Neurosci Abstr* 1999;25:14.

[15] Butterfield DA, Martin L, Carney JM, Hensley K. A β (25–35) peptide displays H₂O₂-like reactivity towards aqueous Fe²⁺, nitroxide spin probes, and synaptosomal membrane proteins. *Life Sci* 1996;58:217–28.

[16] Buxbaum JD, Oishi M, Chen H. Cholinergic agonists and interleukin-1 regulate processing and secretion of the Alzheimer β /A4 amyloid precursor protein. *Proc Natl Acad Sci USA* 1992;89:10075–8.

[17] Buxbaum JD, Liu KN, Luo Y, Slack JL, Stocking KL, Peschon JJ, et al. Evidence that tumor necrosis factor alpha converting enzyme is involved in regulated alpha-secretase cleavage of the Alzheimer amyloid protein precursor. *J Biol Chem* 1998;273:27765–7.

[18] Cai X-D, Golde TE, Younkin SG. Release of excess amyloid- β protein from a mutant amyloid- β protein precursor. *Science* 1993;259:514–6.

[19] Caporaso GL, Gandy SE, Buxbaum JD, Ramabhadran TV, Greengard P. Protein phosphorylation regulates secretion of Alzheimer beta/A4 amyloid precursor protein. *Proc Natl Acad Sci USA* 1992;89:3055–9.

- [20] Carpenter MK, Crutcher KA, Kater SB. An analysis of the effects of Alzheimer's plaques on living neurons. *Neurobiol Aging* 1993;14:207–15.
- [21] Chan C-W, Dharmarajan A, Atwood CS, Huang X, Tanzi RE, Bush AI, et al. Anti-apoptotic action of Alzheimer A β . *Alzheimer's Rep* 1999;2:1–6.
- [22] Ciallella JR, Rangnekar VV, McGillis JP. Heat shock alters Alzheimer's beta amyloid precursor protein expression in human endothelial cells. *J Neurosci Res* 1994;37:769–76.
- [23] Citron M, Oltersdorf T, Haass C, McConlogue L, Hung AY, Seubert P, et al. Mutation of the beta-amyloid precursor protein in familial Alzheimer's disease increases beta-protein production. *Nature* 1992;360:672–4.
- [24] Citron M, Teplow DB, Selkoe DJ. Generation of amyloid-beta protein from its precursor is sequence specific. *Neuron* 1995;14:661–70.
- [25] Citron M, Westaway D, Xia W, Carlson G, Diehl T, Levesque G, et al. Mutant presenilins of Alzheimer's disease increase production of 42-residue amyloid-beta protein in both transfected cells and transgenic mice. *Nat Med* 1997;3:67–72.
- [26] Corder EH, Saunders AM, Strittmatter WJ, Schmechel DE, Gaskell PC, Small GW, et al. Gene dose of apolipoprotein E type 4 allele and the risk of Alzheimer's disease in late onset families. *Science* 1993;261:921–3.
- [27] Corder EH, Saunders AM, Risch NJ, Strittmatter WJ, Roses AD, Haines JL, et al. Protective effect of apolipoprotein E type 2 allele for late onset Alzheimer disease. *Nat Genet* 1994;7:180–4.
- [28] Cotman CW, Pike CJ, Copani A. β -Amyloid neurotoxicity: a discussion of in vitro findings. *Neurobiol Aging* 1992;13:587–90.
- [29] Cuajungco MP, Goldstein LE, Nunomura A, Smith MA, Lim JT, Atwood CS, et al. Evidence that the β -amyloid plaques of Alzheimer's disease represent the redox-silencing and entombment of A β by zinc. *J Biol Chem* 2000;275:19439–42.
- [30] Davies L, Wolska B, Hilbich C, Multhaup G, Martins R, Simms G, et al. A4 amyloid protein deposition and the diagnosis of Alzheimer's disease: prevalence in aged brains determined by immunocytochemistry compared with conventional neuropathologic techniques. *Neurology* 1988;38:1688–93.
- [31] Dikalov SI, Vitek MP, Maples KR, Mason RP. Amyloid- β -peptides do not form peptide-derived free radicals spontaneously, but enhance metal-catalyzed oxidation of hydroxylamines to nitroxides. *J Biol Chem* 1999;274:9392–9.
- [32] Frederikse PH, Garland D, Zigler Jr JS, Piatigorsky J. Oxidative stress increases production of beta-amyloid precursor protein and beta-amyloid (A β) in mammalian lenses, and A β has toxic effects on lens epithelial cells. *J Biol Chem* 1996;271:10169–74.
- [33] Friedman G, Froom P, Sazbon L, Grinblatt I, Shochina M, Tsenter J, et al. Apolipoprotein E- ϵ 4 genotype predicts a poor outcome in survivors of traumatic brain injury. *Neurology* 1999;52:244–8.
- [34] Gabuzda D, Busciglio J, Chen LB, Matsudaira P, Yankner BA. Inhibition of energy metabolism alters the processing of amyloid precursor protein and induces a potentially amyloidogenic derivative. *J Biol Chem* 1994;269:13623–8.
- [35] Games D, Adams D, Alessandrini R, Barbour R, Berthelette P, Blackwell C, et al. Alzheimer-type neuropathology in transgenic mice overexpressing V717F β -amyloid precursor protein. *Nature* 1995;373:523–7.
- [36] Geddes JW, Anderson KJ, Cotman CW. Senile plaques as aberrant sprout-stimulating structures. *Exp Neurol* 1986;94:767–76.
- [37] Geddes JF, Vowles GH, Beer TW, Ellison DW. The diagnosis of diffuse axonal injury: implications for forensic practice. *Neuropathol Appl Neurobiol* 1997;23:339–47.
- [38] Gentleman SM, Nash MJ, Sweeting CJ, Graham DI, Roberts GW. β -Amyloid precursor protein (β APP) as a marker for axonal injury after head injury. *Neurosci Lett* 1993;160:139–44.
- [39] Glenner GG, Wong CW. Alzheimer's disease: initial report of the purification and characterization of a novel cerebrovascular amyloid protein. *Biochem Biophys Res Commun* 1984;120:885–90.
- [40] Glenner GG, Wong CW. Alzheimer's disease and Down's syndrome: sharing of a unique cerebrovascular amyloid fibril protein. *Biochem Biophys Res Commun* 1984;122:1131–5.
- [41] Goate A, Chartier-Harlin MC, Mullan M, Brown J, Crawford F, Fidani L, et al. Segregation of a missense mutation in the amyloid precursor protein gene with familial Alzheimer's disease. *Nature* 1991;349:704–6.
- [42] Goldgaber D, Harris HW, Hla T, Maciag T, Donnelly RJ, Jacobsen JS, et al. Interleukin 1 regulates synthesis of amyloid-beta protein precursor mRNA in human endothelial cells. *Proc Natl Acad Sci USA* 1995;86:7606–10.
- [43] Gooch MD, Stennett DJ. Molecular basis of Alzheimer's disease. *Am J Health Syst Pharm* 1996;53:1545–57.
- [44] Greenberg BD, Savage MJ, Howland DS, Ali SM, Siedlak SL, Perry G, et al. APP transgenesis: approaches toward the development of animal models for Alzheimer disease neuropathology. *Neurobiol Aging* 1996;17:153–71.
- [45] Haass C, Schlossmacher MG, Hung AY, Vigo-Pelfrey C, Mellon A, Ostaszewski BL, et al. Amyloid- β -peptide is produced by cultured cells during normal metabolism. *Nature* 1992;359:322–5.
- [46] Hall ED, Oostveen JA, Dunn E, Carter DB. Increased amyloid protein precursor and apolipoprotein E immunoreactivity in the selectively vulnerable hippocampus following transient forebrain ischemia in gerbils. *Exp Neurol* 1995;135:17–27.
- [47] Harris PLR, Zhu X, Pamies C, Rottkamp CA, Ghanbari HA, McShea A, et al. Neuronal polo-like kinase in Alzheimer disease indicates cell cycle changes. *Neurobiol Aging* 2000;21:837–41.
- [48] Hensley K, Carney JM, Mattson MP, Aksenova M, Harris M, Wu JF, et al. A model for beta-amyloid aggregation and neurotoxicity based on free radical generation by the peptide: relevance to Alzheimer disease. *Proc Natl Acad Sci USA* 1994;91:3270–4.
- [49] Higgins GA, Oyler GA, Neve RL, Chen KS, Gage FH. Altered levels of amyloid protein precursor transcripts in the basal forebrain of behaviorally impaired aged rats. *Proc Natl Acad Sci USA* 1990;87:3032–6.
- [50] Hölscher C. Possible causes of Alzheimer's disease: amyloid fragments, free radicals, and calcium homeostasis. *Neurobiol Dis* 1998;5:129–41.
- [51] Horsburgh K, McCarron MO, White F, Nicoll JAR. The role of apolipoprotein E in Alzheimer's disease, acute brain injury and cerebrovascular disease: evidence of common mechanisms and utility of animal models. *Neurobiol Aging* 2000;21:245–55.
- [52] Hsiao K, Chapman P, Nilsen S, Eckman C, Harigaya Y, Younkin S, et al. Correlative memory deficits, A β elevation, and amyloid plaques in transgenic mice. *Science* 1996;274:99–102.
- [53] Huang X, Atwood CS, Moir RD, Hartshorn MA, Vonsattel JP, Tanzi RE, et al. Zinc-induced Alzheimer's A β _{1–40} aggregation is mediated by conformational factors. *J Biol Chem* 1997;272:26464–70.
- [54] Huang X, Atwood CS, Hartshorn MA, Multhaup G, Goldstein LE, Scarpa RC, et al. The A β peptide of Alzheimer's disease directly produces hydrogen peroxide through metal ion reduction. *Biochemistry* 1999;38:7609–16.
- [55] Huang X, Cuajungco MP, Atwood CS, Hartshorn MA, Tyndall J, Hanson GR, et al. Alzheimer A β interaction with Cu(II) induces neurotoxicity, radicalization, metal reduction and hydrogen peroxide formation. *J Biol Chem* 1999;274:37111–6.
- [56] Jarrett JT, Lansbury Jr PT. Seeding "one-dimensional crystallization" of amyloid: a pathogenic mechanism in Alzheimer's disease and scrapie? *Cell* 1993;73:1055–8.
- [57] Jendroska K, Poewe W, Daniel SE, Pluess J, Iwerssen-Schmidt H, Paulsen J, et al. Ischemic stress induces deposition of amyloid-beta immunoreactivity in human brain. *Acta Neuropathol* 1995;90:461–6.
- [58] Joseph J, Shukitt-Hale B, Denisova NA, Martin A, Perry G, Smith MA. Copernicus revisited: amyloid-beta in Alzheimer's disease. *Neurobiol Aging* 2001;22:131–46.
- [59] Joseph JA, Perry G, Shukitt-Hale B, Denisova NA, Martin A, Smith MA. A diet at amyloid-beta? *Neurobiol Aging* 2001;22:161–3.

- [60] Kaltschmidt B, Uherek M, Wellmann H, Volk B, Kaltschmidt C. Inhibition of NF-kappaB potentiates amyloid-beta-mediated neuronal apoptosis. *Proc Natl Acad Sci USA* 1999;96:9409–14.
- [61] Kang J, Lemaire HG, Unterbeck A, Salbaum JM, Masters CL, Grzeschik KH, et al. The precursor of Alzheimer's disease amyloid A4 protein resembles a cell-surface receptor. *Nature* 1987;325:733–6.
- [62] Kontush A, Berndt C, Arlt S, Schippling S, Beisiegel U. Amyloid- β is a physiological anti-oxidant for lipoproteins in cerebrospinal fluid and plasma. *Neurobiol Aging* 1998;19:43.
- [63] Kontush A, Berndt C, Weber W, Akopyan V, Arlt S, Schippling S, et al. Amyloid-beta is an anti-oxidant for lipoproteins in cerebrospinal fluid and plasma. *Free Radic Biol Med* 2001;30:119–28.
- [64] Koo EH, Park L, Selkoe DJ. Amyloid-beta protein as a substrate interacts with extracellular matrix to promote neurite outgrowth. *Proc Natl Acad Sci USA* 1993;90:4748–52.
- [65] Kovacs DM, Fausett HJ, Page KJ, Kim TW, Moir RD, Merriam DE, et al. Alzheimer-associated presenilins 1 and 2: neuronal expression in brain and localization to intracellular membranes in mammalian cells. *Nat Med* 1996;2:224–9.
- [66] Kowall N, McKee A, Yankner B, Beal M. In vivo neurotoxicity of β -amyloid [β (1–40)] and the β (25–35) fragment. *Neurobiol Aging* 1992;13:537–42.
- [67] Kuentzel SL, Ali SM, Altman RA, Greenberg BD, Raub TJ. The Alzheimer β -amyloid protein precursor/protease nexin-II is cleaved by secretase in a *trans*-Golgi secretory compartment in human neuroglioma cells. *Biochem J* 1993;295:367–78.
- [68] LaFerla FM, Tinkle BT, Bieberich CJ, Haudenschild CC, Jay G. The Alzheimer's A β peptide induces neurodegeneration and apoptotic cell death in transgenic mice. *Nat Genet* 1995;9:21–30.
- [69] Lambert MP, Barlow AK, Chromy BA, Edwards C, Freed R, Liosatos M, et al. Diffusible, nonfibrillar ligands derived from A β _{1–42} are potent central nervous system neurotoxins. *Proc Natl Acad Sci USA* 1998;95:6448–53.
- [70] Lendon CL, Ashall F, Goate AM. Exploring the etiology of Alzheimer disease using molecular genetics. *JAMA* 1997;277:825–31.
- [71] Lorenzo A, Yankner BA. Amyloid fibril toxicity in Alzheimer's disease and diabetes. *Ann NY Acad Sci* 1996;777:89–95.
- [72] Lue LF, Kuo YM, Roher AE, Brachova L, Shen Y, Sue L, et al. Soluble amyloid-beta peptide concentration as a predictor of synaptic change in Alzheimer's disease. *Am J Pathol* 1999;155:853–62.
- [73] Luo Y, Sunderland T, Roth GS, Wolozin B. Physiological levels of beta-amyloid peptide promote PC12 cell proliferation. *Neurosci Lett* 1996;217:125–8.
- [74] Mann DM, Jones D, South PW, Snowden JS, Neary D. Deposition of amyloid-beta protein in non-Alzheimer dementias: evidence for a neuronal origin of parenchymal deposits of beta protein in neurodegenerative disease. *Acta Neuropathol* 1992;83:415–9.
- [75] Masters CL, Simms G, Weinman NA, Multhaup G, McDonald BL, Beyreuther K. Amyloid plaque core protein in Alzheimer disease and Down syndrome. *Proc Natl Acad Sci USA* 1985;82:4245–9.
- [76] Mattson MP. Degenerative and protective signaling mechanisms in the neurofibrillary pathology of AD. *Neurobiol Aging* 1995;16:447–57.
- [77] Mattson MP, Pedersen WA. Effects of amyloid precursor protein derivatives and oxidative stress on basal forebrain cholinergic systems in Alzheimer's disease. *Int J Dev Neurosci* 1998;16:737–53.
- [78] McShea A, Harris PLR, Webster KR, Wahl AF, Smith MA. Abnormal expression of the cell cycle regulators p16 and CDK4 in Alzheimer's disease. *Am J Pathol* 1997;150:1933–9.
- [79] McShea A, Wahl AF, Smith MA. Re-entry into the cell cycle: a mechanism for neurodegeneration in Alzheimer disease. *Med Hypotheses* 1999;52:525–7.
- [80] McShea A, Zelasko DA, Gerst JL, Smith MA. Signal transduction abnormalities in Alzheimer's disease: evidence of a pathogenic stimuli. *Brain Res* 1999;815:237–42.
- [81] Meda L, Cassatella MA, Szendrei GI, Ottovs Jr L, Baron P, Villalba M, et al. Activation of microglial cells by beta-amyloid protein and interferon-gamma. *Nature* 1995;374:647–50.
- [82] Misonou H, Morishima-Kawashima M, Ihara Y. Oxidative stress induces intracellular accumulation of amyloid-beta protein (A β) in human neuroblastoma cells. *Biochemistry* 2000;39:6951–9.
- [83] Murakami N, Yamaki T, Iwamoto Y, Sakakibara T, Kobori N, Fushiki S, et al. Experimental brain injury induces expression of amyloid precursor protein, which may be related to neuronal loss in the hippocampus. *J Neurotrauma* 1998;15:993–1003.
- [84] Nagy Z, Esiri MM, Cato AM, Smith AD. Cell cycle markers in the hippocampus in Alzheimer's disease. *Acta Neuropathol* 1997;94:6–15.
- [85] Nagy Z, Esiri MM, Smith AD. Expression of cell division markers in the hippocampus in Alzheimer's disease and other neurodegenerative conditions. *Acta Neuropathol* 1997;93:294–300.
- [86] Neve RL, Robakis NK. Alzheimer's disease: a re-examination of the amyloid hypothesis. *Trends Neurosci* 1998;21:15–9.
- [87] Nishimoto I, Okamoto T, Matsuura Y, Takahashi S, Okamoto T, Murayama Y, et al. Alzheimer amyloid protein precursor complexes with brain GTP-binding protein G₀. *Nature* 1993;362:14–5.
- [88] Nordstedt C, Gandy SE, Alafuzoff I, Caporaso GL, Iverfeldt K, Grebb JA, et al. Alzheimer beta/A4 amyloid precursor protein in human brain: aging-associated increases in holoprotein and in a proteolytic fragment. *Proc Natl Acad Sci USA* 1991;88:8910–4.
- [89] Nunomura A, Perry G, Pappolla MA, Wade R, Hirai K, Chiba S, et al. RNA oxidation is a prominent feature of vulnerable neurons in Alzheimer's disease. *J Neurosci* 1999;19:1959–64.
- [90] Nunomura A, Perry G, Pappolla MA, Friedland RP, Hirai K, Chiba S, et al. Neuronal oxidative stress precedes amyloid- β deposition in Down syndrome. *J Neuropathol Exp Neurol* 2000;59:1011–7.
- [91] Nunomura A, Perry G, Aliev G, Hirai K, Takeda A, Balraj EK, Jones PK, Ghanbari H, Wataya T, Shimohama S, Chiba S, Atwood CS, Petersen RB, Smith MA. Oxidative damage is the earliest event in Alzheimer disease. *J Neuropathol Exp Neurol* 2001;60:759–67.
- [92] Panegyres PK. The effects of excitotoxicity on the expression of the amyloid precursor protein gene in the brain and its modulation by neuroprotective agents. *J Neural Transmission* 1998;105:463–78.
- [93] Paola D, Domenicotti C, Nitti M, Vitali A, Borghi R, Cottalasso D, et al. Oxidative stress induces increase in intracellular amyloid-beta protein production and selective activation of β I and β II PKCs in NT2 cells. *Biochem Biophys Res Commun* 2000;268:642–6.
- [94] Perry G, Nunomura A, Raina AK, Smith MA. Amyloid- β junkies. *Lancet* 2000;355:757.
- [95] Pike CJ, Walencewicz AJ, Glabe CG, Cotman CW. Aggregation-related toxicity of synthetic beta-amyloid protein in hippocampal cultures. *Eur J Pharmacol* 1991;207:367–8.
- [96] Pike CJ, Cummings BJ, Cotman CW. β -amyloid induces neuritic dystrophy in vitro: similarities with Alzheimer pathology. *Neuroreport* 1992;3:769–72.
- [97] Pike CJ, Burdick D, Walencewicz AJ, Glabe CG, Cotman CW. Neurodegeneration induced by β -amyloid peptide in vitro: role of peptide assembly state. *J Neurosci* 1993;13:1676–87.
- [98] Pike CJ, Walencewicz-Wasserman AJ, Kosmoski J, Cribbs DH, Glabe CG, Cotman CW. Structure-activity analyses of beta-amyloid peptides: contributions of the beta 25–35 region to aggregation and neurotoxicity. *J Neurochem* 1995;64:253–65.
- [99] Poirier J, Baccichet A, Dea D, Gauthier S. Cholesterol synthesis and lipoprotein reuptake during synaptic remodeling in hippocampus in adult rats. *Neuroscience* 1993;55:81–90.
- [100] Postuma RB, He W, Nunan J, Beyreuther K, Masters CL, Barrow CJ, et al. Substrate-bound beta-amyloid peptides inhibit cell adhesion

- and neurite outgrowth in primary neuronal cultures. *J Neurochem* 2000;74:1122–30.
- [101] Raina AK, Monteiro MJ, McShea A, Smith MA. The role of cell cycle-mediated events in Alzheimer's disease. *Int J Exp Pathol* 1999;80:71–6.
- [102] Raina AK, Takeda A, Smith MA. Mitotic neurons: a dogma succumbs. *Exp Neurol* 1999;159:248–9.
- [103] Raina AK, Zhu X, Monteiro M, Takeda A, Smith MA. Abortive oncogeny and cell cycle-mediated events in Alzheimer disease. In: Meijer L, Jézéquel A, Ducommun B, editors. *Progress in cell cycle research*, vol. 4. New York: Kluwer Academic Publishers/Plenum Press, 2000. p. 235–42.
- [104] Raina AK, Zhu X, Rottkamp CA, Monteiro M, Takeda A, Smith MA. Cyclin' toward dementia: cell cycle abnormalities and abortive oncogenesis in Alzheimer disease. *J Neurosci Res* 2000;61:128–33.
- [105] Raina AK, Hochman A, Zhu X, Rottkamp CA, Nunomura A, Siedlak SL, et al. Abortive apoptosis in Alzheimer's disease. *Acta Neuropathol* 2001;101:305–10.
- [106] Roberts GW, Gentleman SM, Lynch A, Murray L, Landon M, Graham DI. Beta amyloid protein deposition in the brain after severe head injury: implications for the pathogenesis of Alzheimer's disease. *J Neurol Neurosurg Psychiatr* 1994;57:419–25.
- [107] Rogers J, Morrison JH. Quantitative morphology and regional and laminar distributions of senile plaques in Alzheimer's disease. *J Neurosci* 1985;5:2801–8.
- [108] Rottkamp CA, Nunomura A, Raina AK, Sayre LM, Perry G, Smith MA. Oxidative stress, anti-oxidants, and Alzheimer disease. *Alzheimer Dis Assoc Disord* 2000;14(Suppl 1):S62–6.
- [109] Rottkamp CA, Raina AK, Zhu X, Gaier E, Bush AI, Atwood CS, et al. Redox-active iron mediates amyloid- β toxicity. *Free Radic Biol Med* 2001;30:447–50.
- [110] Sayre LM, Zagorski MG, Surewicz WK, Krafft GA, Perry G. Mechanisms of neurotoxicity associated with amyloid- β deposition and the role of free radicals in the pathogenesis of Alzheimer's disease: a critical appraisal. *Chem Res Toxicol* 1997;10:518–26.
- [111] Sayre LM, Perry G, Harris PLR, Liu Y, Schubert KA, Smith MA. In situ oxidative catalysis by neurofibrillary tangles and senile plaques in Alzheimer's disease: a central role for bound transition metals. *J Neurochem* 2000;74:270–9.
- [112] Schenk D, Barbour R, Dunn W, Gordon G, Grajeda H, Guido T, et al. Immunization with amyloid-beta attenuates Alzheimer-disease-like pathology in the PDAPP mouse. *Nature* 1999;400:173–7.
- [113] Selkoe DJ. Amyloid- β -protein and genetics of Alzheimer's disease. *J Biol Chem* 1996;271:18295–8.
- [114] Selkoe DJ. Translating cell biology into therapeutic advances in Alzheimer's disease. *Nature* 1999;399(suppl):A23–31.
- [115] Shi J, Xiang Y, Simpkins JW. Hypoglycemia enhances the expression of mRNA encoding beta-amyloid precursor protein in rat primary cortical astroglial cells. *Brain Res* 1997;772:247–51.
- [116] Shi J, Panickar KS, Yang SH, Rabbani O, Day AL, Simpkins JW. Estrogen attenuates over-expression of beta-amyloid precursor protein messenger RNA in an animal model of focal ischemia. *Brain Res* 1998;810:87–92.
- [117] Shi J, Yang SH, Stubble L, Day AL, Simpkins JW. Hypoperfusion induces overexpression of beta-amyloid precursor protein mRNA in a focal ischemic rodent model. *Brain Res* 2000;853:1–4.
- [118] Shoji M, Golde TE, Ghiso J, Cheung TT, Estus S, Shaffer LM, et al. Production of the Alzheimer amyloid-beta protein by normal proteolytic processing. *Science* 1992;258:126–9.
- [119] Singh VK, Cheng JF, Leu SJ. Effect of substance P and protein kinase inhibitors on beta-amyloid peptide-induced proliferation of cultured brain cells. *Brain Res* 1994;660:353–6.
- [120] Sisodia SS. Secretion of the beta-amyloid precursor protein. *Ann NY Acad Sci* 1992;674:53–7.
- [121] Smith TW, Lipp CF. Ki-67 immunoreactivity in Alzheimer's disease and other neurodegenerative disorders. *J Neuropathol Exp Neurol* 1995;54:297–303.
- [122] Smith MA, Taneda S, Richey PL, Miyata S, Yan SD, Stern D, et al. Advanced Maillard reaction products are associated with Alzheimer disease pathology. *Proc Natl Acad Sci USA* 1994;91:5710–4.
- [123] Smith MA, Harris PLR, Sayre LM, Perry G. Iron accumulation in Alzheimer disease is a source of redox-generated free radicals. *Proc Natl Acad Sci USA* 1997;94:9866–8.
- [124] Smith MA, Hirai K, Hsiao K, Pappolla MA, Harris PLR, Siedlak SL, et al. Amyloid- β deposition in Alzheimer transgenic mice is associated with oxidative stress. *J Neurochem* 1998;70:2212–5.
- [125] Smith MA, Joseph JA, Perry G. Arson: tracking the culprit in Alzheimer's disease. *Ann NY Acad Sci* 2000;924:35–8.
- [126] Sorbi S, Nacmias N, Piacentini S, Repice A, Latorraca S, Forleo P, et al. ApoE as a prognostic factor for post-traumatic coma. *Nat Med* 1995;1:852.
- [127] Stephenson DT, Rash K, Clemens JA. Amyloid precursor protein accumulates in regions of neurodegeneration following focal cerebral ischemia in the rat. *Brain Res* 1992;593:128–35.
- [128] Suzuki N, Cheung T, Cai X, Odaka A, Ötvös LJ, Eckman C, et al. An increased percentage of long amyloid-beta protein secreted by familial amyloid-beta protein precursor (BAPP717) mutants. *Science* 1994;264:1336–40.
- [129] Takenouchi T, Munekata E. Trophic effects of substance P and beta-amyloid peptide on dibutyryl cyclic AMP-differentiated human leukemic (HL-60) cells. *Life Sci* 1995;56:PL479–84.
- [130] Teller JK, Russo C, DeBusk LM, Angelini G, Zaccheo D, Dagna-Bricarelli F, et al. Presence of soluble amyloid-beta peptide precedes amyloid plaque formation in Down's syndrome. *Nat Med* 1996;1:93–5.
- [131] Tomita T, Maruyama K, Saido TC, Kume H, Shinozaki K, Tokuhira S, et al. The presenilin 2 mutation (N141I) linked to familial Alzheimer disease (Volga German families) increases the secretion of amyloid-beta protein ending at the 42nd (or 43rd) residue. *Proc Natl Acad Sci USA* 1997;94:2025–30.
- [132] Topper R, Gehrman J, Banati R, Schwarz M, Block F, Noth J, et al. Rapid appearance of beta-amyloid precursor protein immunoreactivity in glial cells following excitotoxic brain injury. *Acta Neuropathol* 1995;89:23–8.
- [133] Turnbull S, Tabner BJ, El-Agnaf OMA, Twyman LJ, Allsop D. New evidence that the Alzheimer β -amyloid peptide does not spontaneously form free radicals: an ESR study using a series of spin-traps. *Free Radic Biol Med* 2001;30:1154–62.
- [134] van Gool WA, Schenk DB, Bolhuis PA. Concentrations of amyloid-beta protein in cerebrospinal fluid increase with age in patients free from neurodegenerative disease. *Neurosci Lett* 1994;172:122–4.
- [135] Van Muiswinkel FL, Veerhuis R, Eikelenboom P. Amyloid-beta protein primes cultured rat microglial cells for an enhanced phorbol12-myristate 13-acetate-induced respiratory burst activity. *J Neurochem* 1996;66:2468–76.
- [136] Vincent I, Rosado M, Davies P. Mitotic mechanisms in Alzheimer's disease? *J Cell Biol* 1996;132:413–25.
- [137] Vincent I, Jicha G, Rosado M, Dickson DW. Aberrant expression of mitotic cdc2/cyclinB1 kinase in degenerating neurons of Alzheimer's disease brains. *J Neurosci* 1997;17:3588–98.
- [138] Wallace W, Ahlers ST, Gotlib J, Bragin V, Sugar J, Gluck R, et al. Amyloid precursor protein in the cerebral cortex is rapidly and persistently induced by loss of subcortical innervation. *Proc Natl Acad Sci USA* 1993;90:8712–6.
- [139] Whitson JS, Selkoe DJ, Cotman CW. Amyloid-beta protein enhances the survival of hippocampal neurons in vitro. *Science* 1989;243:1488–90.
- [140] Whitson JS, Glabe CG, Shintani E, Abcar A, Cotman CW. Beta-amyloid protein promotes neuritic branching in hippocampal cultures. *Neurosci Lett* 1990;110:319–24.
- [141] Wisniewski T, Castano EM, Golabek A, Vogel T, Frangione B. Acceleration of Alzheimer's fibril formation by apolipoprotein E in vitro. *Am J Pathol* 1994;145:1030–5.

- [142] Xia W, Zhang J, Perez R, Koo EH, Selkoe DJ. Interaction between amyloid precursor protein and presenilins in mammalian cells: implications for the pathogenesis of Alzheimer disease. *Proc Natl Acad Sci USA* 1997;94:8208–13.
- [143] Xiong Y, Gu Q, Peterson PL, Muizelaar JP, Lee CP. Mitochondrial dysfunction and calcium perturbation induced by traumatic brain injury. *J Neurotrauma* 1997;14:23–34.
- [144] Yan SD, Chen X, Schmidt AM, Brett J, Godman G, Zou YS, et al. Glycated tau protein in Alzheimer disease: a mechanism for induction of oxidant stress. *Proc Natl Acad Sci USA* 1994;91:7787–91.
- [145] Yankner BA, Duffy LK, Kirschner DA. Neurotrophic and neurotoxic effects of amyloid-beta protein: reversal by tachykinin neuropeptides. *Science* 1990;250:279–82.
- [146] Yankner BA. New clues to Alzheimer's disease: unraveling the roles of amyloid and tau. *Nature Med* 1996;2:850–2.
- [147] Yokota M, Saido TC, Tani E, Yamaura I, Minami N. Cytotoxic fragment of amyloid precursor protein accumulates in hippocampus after global forebrain ischemia. *J Cerebral Blood Flow Metab* 1996;16:1219–23.
- [148] Zhu X, Raina AK, Smith MA. Cell cycle events in neurons: proliferation or death? *Am J Pathol* 1999;155:327–9.
- [149] Zhu X, Raina AK, Boux H, Simmons ZL, Takeda A, Smith MA. Activation of oncogenic pathways in degenerating neurons in Alzheimer disease. *Int J Dev Neurosci* 2000;18:433–7.
- [150] Zhu X, Rottkamp CA, Boux H, Takeda A, Perry G, Smith MA. Activation of p38 kinase links tau phosphorylation, oxidative stress and cell cycle-related events in Alzheimer disease. *J Neuropathol Exp Neurol* 2000;59:880–8.
- [151] Zhu X, Rottkamp CA, Raina AK, Brewer GJ, Ghanbari HA, Boux H, et al. Neuronal CDK7 in hippocampus is related to aging and Alzheimer disease. *Neurobiol Aging* 2000;21:807–13.
- [152] Bishop GM, Robinson SR, Smith MA, Perry G, Atwood CS. Call for Elan to publish Alzheimer's trial details (Letter). *Nature* 2002;416:677.
- [153] Check E. Nerve inflammation halts trial for Alzheimer's drug (News). *Nature* 2002;415:462.
- [154] Steinberg D. Companies halt first Alzheimer vaccine trial. *The Scientist* 2002;16:22–3.
- [155] Weiss R. Alzheimer's Vaccine Permanently Shelved. Testing's Toll of Brain Ailments Grows. *Washington Post*, 2 March 2002, A03.